

Discharge Instructions Facelift/Necklift/Fat Transfer

Dr Sean Blaydon Dr. Tanuj Nakra Dr. Vikram Durairaj Dr. Irina Belinsky Dr. Donovan Reed
 Dr. Anish Abrol Dr. Monica Ray Dr. Mark Mazow Dr. Karen Brown

After hours & weekends (512) 533-7355
Office (512) 458-2141 M-F 8 am-5pm

CALL IMMEDIATELY FOR THESE PROBLEMS:

- Pain not controlled by medication.
- Fever over 101 degrees with increasing pain, redness, swelling, and drainage from wound.
- If you notice rapid swelling or blood dripping from the incision.

DRESSING

- Keep the facial/neck dressing dry, and leave dressing in place until seen by your surgeon in follow-up.
- The dressing on the fat transfer donor site may be removed the next day. Expect drainage from the donor site for first 24 hrs.

MEDICATION

- Continue your regular medications unless otherwise instructed by physician.
- Apply a small amount of _____ ointment to the stitches _____ times a day for _____ days.
- Take prescription **pain** medication for moderate pain. Tylenol/Extra Strength Tylenol for mild pain. Take pain medication with food to avoid nausea.
- You were given pain medication in the surgery center. Next dose of pain medication at _____.*
- DO NOT** take Aspirin, Aspirin containing products or anti-inflammatory medications (NSAIDS e.g. Advil, Motrin, Aleve, Celebrex, etc.), for 7 days after surgery.
- Blood thinning medication such as Coumadin, Plavix or Aspirin can be restarted on _____.*
- Other medications: Promethazine as needed for nausea. Next dose: _____
 Medrol Dose Pack starts _____
- Rest quietly at home for two days after surgery and elevate your head on 2-3 pillows while sleeping.
- Stay in the company of a responsible adult for 24 hours after surgery.
- When bending, bend at knees only, and not from waist.
- Avoid lifting anything heavier than a gallon of milk for 7 days.
- No heavy exercising of any kind for two to three weeks, and until approved by your surgeon.
- Wear a cervical foam collar after the bandage is removed to protect your neck, and remove it only for showering.
- Use caution in turning your head and avoid exaggerated head movements for two weeks. Drive only in familiar areas.
- Wear button down shirts to minimize use of pullover clothing for 2 weeks.
- Do not submerge your head underwater in a swimming pool or hot tub for 1 month after surgery.
- **Do not drive** while taking prescription pain medication. **Do not smoke** for 2 – 4 weeks following surgery.

HYGIENE

- After the dressings are removed you may shower and gently shampoo your hair.
- Numbness of the ears and neck may be present for some weeks after surgery. If using a hair dryer, use with caution on warm setting. Avoid prolonged exposure to cold temperatures.

DIET

- Resume regular diet. Select foods that are easily chewed and take small bites.
- No alcoholic beverages for 1 week after surgery, or while taking prescription pain medication.

OTHER _____

FOLLOW UP APPOINTMENT _____

If you have questions immediately after surgery, call Dr. _____ Mon-Fri 8am-5pm at 512-458-2141