	Discharge Instructions Rhinoplasty	Date:
CALL Pa Fe	aydon Dr. Tanuj Nakra Dr. Vikram Durairaj n Reed Dr. Monica Ray Dr. Mark Mazow IMMEDIATELY FOR THESE PROBLEMS: in not controlled by medication. ver over 101 degrees with increasing pain, redness, sween see bleed that is not controlled with pressure for 10 minutes.	
DRESSING Do	not remove the splint on your nose or get it wet.	
	ange the drip pad under your nose as needed for the next	24-48 hours.
MEDICATIO		11 1 2 2
	ntinue your regular medications unless otherwise instruct	
	Apply a small amount of Neosporin ointment to the stitches 2 times a day for 7 days. Take prescription pain medication for moderate pain. Tylenol/Extra Strength Tylenol for mild pain.	
	Take pain medication with food to avoid nausea.	
	You were given pain medication in the surgery center. Next dose of pain medication at	
	Do not take Aspirin or anti-inflammatory medications (Advil, Motrin, Aleve, etc.), for 2 days after surgery.	
☐ Oth	er medications: Ondansetron as needed for nausea. No	
	☐ Medrol Dose Pack starts	
		sture; Starting Day 3, spray saline into nostrils
ACTIVITY	and gently blow into sink.	
	eep on your back or side with your head elevated on 2-3 p	pillows
	by in the company of a responsible adult for 24 hours after	
	When bending, bend at knees only, and not from waist.	
	Avoid lifting anything heavier than a gallon of milk for 7 days.	
• Do	Do not submerge your head underwater in a swimming pool or hot tub for 1 month after surgery.	
• Do	not smoke for 2-4 weeks following your surgery.	
	not drive while taking prescription pain medication.	
HYGIENE		
	ou may shower the day after surgery, but do not get your	<u>-</u>
	e Q-tip dipped in Hydrogen Peroxide to gently clean just	t inside the nostrils.
DIET		
	sume regular diet. No alcoholic beverages for 2 days after the plenty of fluids. For the first 48 hours, cool or lukew	
OTHER		
FOLLOW UP	APPOINTMENT	
If you have qu	estions immediately after surgery, call Dr	Mon-Fri 8am-5pm at 512-458-2141